

Symptoms of cranio-mandible dysfunction or bruxism among students

(Objawy dysfunkcji czaszkowo- żuchwowej lub bruksizmu wśród studentów)

M Dołoszycka ^{1,A,D}, M Kulesa- Mrowiecka ^{1,A,E}, Z Kopański ^{1,F}, D Krzemiński ^{2,C,E},
W Ptak ^{2,C}, S Dyl ^{2,B}, I Sklyarov ^{3,B}

Abstract – Introduction. Nowadays, there is a clear increase in the incidence of functional disorders of the stomatognathic system. This is probably related to the growing need to care for your health, but it also may result from a real increase in the incidence of this pathology. High relevance of the issues, prompted the authors to undertake their own research.

Aim of the study. The aim of the study was to assess the frequency of the occurrence of bruxism symptoms among students of Krakow's universities.

Material and methods. The study group consisted of 277 people (172 women and 105 men) aged 19 – 29 who were students of Krakow's universities. An original questionnaire was used and a short physical examination was carried out.

Results and conclusions. The most frequently felt ailment among the subjects was headache. It was reported by 80% of respondents, followed by a disturbed mandibular movement pathway, which was recorded in 78% of respondents, cracks in the temporomandibular joint, which were given by 50% of subjects: presence of tooth clashes and lack of symmetry of masseter muscles, which were found at the same frequency in 38 % of respondents. Tinnitus related to 22.74% of the subjects. A disturbed mandibular disorder pathway occurred in 77.98% of patients with masticatory system dysfunction.

Key words - symptoms of masticatory dysfunction, students.

Streszczenie – Wstęp. Współcześnie odnotowuje się wyraźny wzrost częstości występowania zaburzeń czynnościowych układu stomatognatycznego. Ma to zapewne związek ze wrastającą potrzebą dbałości o swoje zdrowie, ale także może wynikać z rzeczywistego wzrostu częstości występowania tej patologii. Duża aktualność problematyki skłoniła autorów do podjęcia badań własnych.

Cel pracy. Celem badań była ocena częstości występowanie objawów bruksizmu wśród studentów krakowskich uczelni wyższych.

Materiał i metody. Grupę badaną stanowiło 277 osób (172 kobiety i 105 mężczyzn) w wieku 19 - 29 lat, studentów krakowskich uczelni. Wykorzystano autorską ankietę oraz przeprowadzono krótkie badanie fizykalne.

Wyniki i wnioski. Najczęściej odczuwaną dolegliwością wśród badanych był *ból głowy*. Zgłaszało go 80% respondentów, następnie *zaburzony tor ruchu odwodzenia żuchwy*, który rejestrowano u 78% badanych, *trzaski w stawie skroniowo- żuchwowym*, które podawało 50% badanych: *obecność starć zębów i brak symetrii napięcia mięśni żwaczy*, które stwierdzano z jednakową częstością u 38% badanych. Szumy uszne dotyczyły 22,74% badanych. Zaburzony tor odwodzenia żuchwy występował u 77,98% chorych z dysfunkcją narządu żucia .

Wnioski. Jednym z wiodących , choć niecharakterystycznych objawów dysfunkcji narządu żucia jest ból.

Chorym z zaburzeniami układu stomatognatycznego towarzyszyć mogą różnorodne objawy otologiczne występujące w dużym przedziale częstości.

Słowa kluczowe - objawy dysfunkcji narządu żucia, studenci.

Author Affiliations:

1. Faculty of Health Sciences, Collegium Medicum, Jagiellonian University
2. Collegium Masoviense – College of Health Sciences, Żyrardów
3. Department of Therapy and Medical Diagnosis, Faculty of Postgraduate Education, Lviv National Medical University, Ukraine

Authors' contributions to the article:

- A. The idea and the planning of the study
- B. Gathering and listing data
- C. The data analysis and interpretation
- D. Writing the article

E. Critical review of the article
F. Final approval of the article

Correspondence to:

Prof. Zbigniew Kopański MD PhD, Faculty of Health Sciences,
Collegium Medicum, Jagiellonian University, Piotra
Michałowskiego 12 Str., PL- 31-126 Kraków, Poland, e-mail:
zkopanski@o2.pl

Accepted for publication: November 30, 2018.

I. INTRODUCTION

Nowadays, there is a clear increase in the frequency of functional disorders of the stomatognathic system. This is probably related to the growing need to care for your health, but it may also result from a real increase in the incidence of this pathology. [1-7] It is estimated that in the last twenty years in Poland, the number of patients with masticatory dysfunctions doubled. [1]. These dysfunctions occupy the third place in the group of the most common oral pathologies, giving way to caries and periodontal diseases. [4] Depending on the criteria used, it is estimated that approximately 80-80% of adults affect 75% of adults and up to 75% of adolescents aged 15-17 [1.7-9]. The aim of the study was to assess the frequency of the occurrence of bruxism symptoms among students of Kraków's universities.

II. MATERIALS AND METHODS

Materials

The survey was conducted from March 1 to May 31, 2014 in Krakow. The studied group comprised 277 students of four Cracow universities: Collegium Medicum of the Jagiellonian University, AGH University of Science and Technology Stanisława Staszica, Cracow University of Technology Tadeusz Kościuszko and the Pedagogical University named after National Education Commission (Rectors of Cracow universities were obtained for research). Universities were represented in the study respectively by 66 (Collegium Medicum of the Jagiellonian University), 69 (AGH University of Science and Technology), 74 (Cracow University of Technology) and 68 (Pedagogical University) students. 62% (172 people) of the study group were women. The subjects were aged from 19 to 29 years.

Methods

The survey was voluntary, anonymous and random. It used an original questionnaire containing questions about age, sex, year and field of study and questions about the occurrence of physical symptoms such as: headache, jaw pain, cracks in the temporomandibular joint, tinnitus, tooth fracture for an unclear reason, waking up at night. The subject determined the frequency of occurrence of a given symptom.

The questions concerned the occurrence of various parafunctions, for example: chewing gum, chewing pens - the subject marked the frequency of practicing the parafunction. In the question assessing susceptibility to stress, the subject had to mark the answer, which according to him best describes his reactions to stressful situations.

The respondent was also assessed by asking questions whether he was a hyperactive person. As an explanation of this term, it is stated that hyperactivity occurs when a person is unable to sit for one full hour without changing his position or performing additional movements.

Another behavior that was asked in the survey was practicing sports, the respondent reported what sports he was practicing and how many hours in a week he spent on physical activity.

The further part of the questionnaire concerned the method of feeding in early childhood, the researcher indicated whether he was fed naturally (breast) or by a bottle (there was an answer "I do not know" until the election).

Immediately after the survey a physical examination was carried out. The masseter muscle tension on the trailers and belly was checked - the symmetrical pressure of the muscle attachment sites and the muscular thigh was used. The subject valued his feeling and gave it weight in the ten-point VAS scale. The masseter muscle tension was examined using the VAS scale to record the results. For the purpose of the analysis, the examined students were divided according to the reported muscle tone of the masseter into 3 groups: low voltage (1-4 on the VAS scale), medium (5-6) and high (7-10).

The range of abduction of the mandible was measured using a 3-finger test; the examined person checked if he was able to open his mouth to such a width as to fit his own 3 fingers (index, middle and ring finger) between the edges of the lower and upper arch teeth. To ensure hygiene during this test, the patient was instructed to use disinfectant.

Then the subject assessed the occurrence of inequalities and swellings on the mucous membrane of the cheeks. He independently checked the inside of the cheeks with his

tongue, giving the examiner whether these swellings are present.

Another symptom that was checked in the study was hypermobility - passive thumb pull to the forearm was used. The result was considered positive if the subject managed to touch the forearm with the thumb.

The researcher then assessed the tooth abrasion by comparing the height of the upper canines relative to the upper lateral incisors. If on one side there was an alignment of the height of the canine with the side incisor, a positive sign of clash was considered.

The mandibular abduction pathway was also examined, the subject performed several abortions, giving the examiner the opportunity to assess the course of the stereotypical trajectory. Any non-straight track was treated as a disorder.

Often one of the subjects presented several types of the abduction track, which is why the movement was repeated several times, which allowed to clearly determine how the typical diversion track is going for a given patient. The result of this test was marked on the Farrar diagram.

III. RESULTS

The occurrence of symptoms of cranio-mandibular dysfunction or bruxism

The incidence of symptoms of cranio-mandible dysfunction or bruxism among the analyzed group of students is presented in Table 1.

Table 1. The incidence of symptoms of cranio-mandible dysfunction or bruxism in the study group

Symptom		Occurrence
Headache	221	79,78%
Jaw pain felt after waking up	44	15,88%
Cracks in the temporomandibular joint	138	49,82%
Tinnitus	63	22,74%
Hyperactivity	24	8,66%
Breaking a tooth without a clear cause	24	8,66%
Nocturnal awakening	77	27,80%
Hypermobility	100	36,10%
Convergence disorder	44	15,88%
Increased masseter tone: right eye / left eye	96/91	34,66%/32,85%
No symmetry of the masseter muscles tension	105	37,91%
Limited movement of abduction of the mandible	8	2,89%
Disturbed mandibular movement path	216	77,98%
The presence of swellings on the cheeks	97	35,02%
Tooth abrasion	105	37,91%

In the group of symptoms presented in Table 1, there is no difference in the symptoms of craniofacial dysfunction or bruxism, as well as tooth abrasion.

The most felt ailment among the subjects was headache. It was reported by 80% of respondents, which was recorded in 100% of respondents, cracks in the temporo-mandibular

joint; symmetry of masseter muscles, which were found at 38% of respondents.

The incidence of pain in the study group is shown in Table 2

Table 2. Distribution of the occurrence of the study group

Frequency of head-aches	Number	%
everyday	2	0,72
3-5 days in a week	14	5,05
1-2 days in a week	53	19,13
1 in a month	128	46,21
never	56	20,22
other ¹	24	8,66
Altogether	277	100,00

1 / In the "other" category there is a frequency of less than 1 per month, as well as pains that the subjects directly connect to a specific situation (e.g. alcohol consumption), and pain occurring irregularly and sporadically.

The presented data show that headaches in 46.21% of subjects were reported once a month, in 19.13% they occurred with a frequency of 1-2 times a week. Other periodicals of headache were definitely less frequent.

The distribution of the frequency of occurrence of cracks in the temporomandibular joint in the studied group is presented in Table 3.

Table 3. The occurrence of cracks in the temporomandibular joint in the study group

The frequency of occurrence of cracks in the temporomandibular joint	Number	%
often	31	11,19
seldom	107	38,63
never	139	50,18
Altogether	277	100,00

Cracks in the temporomandibular joint occurring during the chewing activity were observed by 49.72% of students, with 11.19% of them finding that they occur frequently.

The distribution of tinnitus frequency in the study group is shown in Table 4.

Table 4. Distribution of occurrence of tinnitus in the study group

Presence of tinnitus	Number	%
every few days	5	1,81
every few hours	3	1,08
several times a week	55	19,86
never	214	77,26
Altogether	77	100,00

22.74% of the surveyed students confirmed tinnitus, 19.86% were of the opinion that these symptoms occur several times a week, other symptoms were much less frequent.

The distribution of nocturnal awakenings in the study group is presented in Table 5.

Table 5. Occurrence of nocturnal awakenings in the study group by frequency

Frequency of awakenings	Number	%
1-2/ night	13	4,69
1/ night	10	3,61
2-5/ week	17	6,14
1/ week	4	1,44
2-4/ month	10	3,61
1-2/ month	23	8,30
none	200	72,20%
Altogether	277	100,00

27.80% of respondents admitted to waking up at night. Nocturnal awakenings occurred most frequently (8.3% of respondents) once or twice in a month, in 6.14%: 2-5 awakenings per week, and in 4.69%: 1-2 every night. The distribution of prevalence of convergence disorders in the study group is presented in Table 6.

Table 6. Occurrence of convergence disorders in a group of students

Convergence disorder	Number	%
convergence is correct	233	84,12
convergence disturbed only in the left eye	25	9,03
convergence disturbed only in the right eye	6	2,17
convergence disturbed in both eyes	13	4,69
Altogether	277	100,00

Convergence disorders affecting at least one eye occurred in 15.88% of subjects, with the most common (9.03% of students) in the left eye.

The distribution of masseter muscle tone in the study group is shown in Table 7.

Table 7. Distribution of masseter muscles tension in the study group

Tension of the masseter muscles	Right		Left	
	Number	%	Number	%
high	34	12,27	27	9,75
average	62	22,38	64	23,10
low	181	65,34	186	67,15
Altogether	277	100,00	277	100,00

The lowest frequency of masseter muscles was recorded in the subjects, 65.34% for the right eye and 67.15% for the left eye. Also, the average muscle tone of the masseter occurred symmetrically with the similar frequency (for the right eye: in 22.38% of cases, for the left eye: in 23.10% of cases). The intensity of masseter muscle tension did not show statistically significant differences between the right and left eyes.

The occurrence of asymmetry in the masseter muscles tension in the examined group is presented in Table 8.

Table 8. The division of students into groups due to the occurrence of asymmetry in the tension of masseter muscles

The occurrence of asymmetry in the strain of masseter muscles	Number	%
Full symmetry	172	62,09
1 or 2	89	32,13
3 or 4	12	4,33
5 or 6	2	0,72
7 and more	2	0,72
Altogether	277	100,00

The asymmetry in masseter muscle tension was noted in 37.91% of the subjects, with asymmetry 1 or 2 being the most frequent in 3213 students.

The restriction of mandibular abduction movement concerns 2.89% of respondents, whereas the disorder of this movement was found in 77.98% of them. Five types of thoracic abduction were distinguished from the results of the tests carried out. The graphical type is presented in Figure 1. One more rare type was observed, which cannot be presented unambiguously, because it is a combination of several arcs in different directions, often with crossing the midline of the body.




Figure 1. Presentations of tracks for the movement of mandibular abduction

Ablasion of fangs was detected in 37.91%, while the presence of swellings and irregularities on the cheeks in 35.02% of the respondents.

Table 9. Prevalence of selected symptoms of craniomandible dysfunction or bruxism in the studies of various authors

Authors	Sójka A., Hędzalek W. [27]	Nawrocka-Furmanek i wsp. [38]	Litko M. Kleinrok J.[1]	Badania własne
Respondent's age	20-25	19-25	Children and teenagers	19-29
The size of the studied group	30	264	Literature review	277
Teeth abrasion	60%	The relationship between the pathological clash and the symptoms of masticatory dysfunction was not found	Not included	38%
Increased muscle tone of the masticatory	76%	The relationship between the back of the bite and the soreness of the cheek	Relationship of muscle tension with execution parafunctions	65%
Acoustic symptoms	56%	More often in the group with nieprawidłowości tooth	Not studied	50% of respondents, more often occur with noise tinnitus
Orthodontic treatment	Not studied	There was no correlation with the occurrence of symptoms of masticatory organ dysfunction	The authors pay attention to the iatrogenic pain syndromes of the masticatory system occurring after orthodontic treatment	46% in past or present. No there were found dependencies with the occurrence of symptoms of masticatory organ dysfunction

IV. DISCUSSION

It is widely emphasized that one of the leading, though non-characteristic symptoms of masticatory system dysfunction is pain. This type of discomfort is referred to as myofascial pain or maxillo-facial pain (maxillofacial pain). Its origin is interpreted differently. It indicates the in-

involvement of impaired functions of the masticatory system, the joint or muscular background of the ailments involved, as well as the involvement of vascular diseases, neurological and psychiatric disorders, and systemic disorders in the formation of this type of discomfort. The pain may be accompanied by local warming, intensifying when changing the position of the masticatory system. Muscle pain is usually constant, dull, diffuse, and increases with increased muscular activity of the masticatory organ. Probably this type of discomfort is a consequence of irritation of nerve endings with local metabolites formed as a result of muscle work and irritation of perivascular plexuses of contracting muscles. [10-21]

Nowadays, each of us is subjected to the activity of numerous stressors. The consequence of emerging stressful situations is the prolonged stimulation of the limbic system and, as a result, the contraction of the masticatory muscles. This promotes self-destruction and the growth of pathology of the masticatory organ. It comes to excessive teeth abrasion, vertical cracking within tooth enamel, lowering of the height of the shorting, hyperfunction of the masticatory muscle muscles. [22-25]

Excessive tooth abrasion is a relatively common symptom of masticatory system dysfunction. According to Maciejewska-Szaniec data, this symptom is accompanied by the dysfunction of the masticatory system from 29.41 to 93.55% of cases, depending on the selection of the analyzed group. [26]

The result of 37.91% obtained by us is therefore, in the cited range. In turn, Sójka and Hędzulek [27] when analyzing a group of 30 students observed that teeth abrasion was experienced by 60% of the respondents. The discrepancy between our data and the results of other authors may result in the adopted assessment criteria. The results obtained by us may also be influenced by the fact that only fangs were checked in our study, while in our assessment, for example, Sójka and Hędzulek considered all teeth.

Patients with disorders of the stomatognathic system may be accompanied by various otological symptoms. These include tinnitus, feeling of blockage, ear and pre-bursal hearing, temporary or even permanent hearing loss. According to literature data, they occur at a frequency of 2.0-82.4%. [28-31]

In Maciejewska-Szaniec studies, otological symptoms were present in 31.69% of patients with masticatory dysfunction [26]. According to our findings, tinnitus refers to 22.74% of respondents. In contrast, Michalak et al. reported otological symptoms in 33.38% [32].

In turn, the aforementioned Sójka and Hędzulek found that the occurrence of acoustic symptoms were identified in

56% of the respondents. The occurrence of tinnitus is variously translated. Attention was paid to the possibility of oppression of the bone elements on the anatomy / temporal nerve, vein and anterior tympanic artery as well as on the external auditory canal and the auditory trumpet, which would cause disturbances of blood supply, innervation and hearing organ function [33,34].

This type of ailment was also associated with direct mechanical stimulation of the hammer, through its connection with the structures of the temporomandibular joint (otomandibular ligate of hypotesis) [35]

Myrhaug emphasized that the common nerves of the muscles of the trigeminal nerve are the basis of aural problems: the tension of the palate, tympanic membrane and masseter muscles. [36]

Ear problems were also associated with the contraction of the blood vessels supplying the ear receptor. Nowadays it is more and more often emphasized that otological complaints in patients with mastectar dysfunction may be associated with irritation of the trigeminal nerve.

According to our findings, the disturbed mandibular abdomen is present in 77.98% of patients with masticatory dysfunction. Sójka and Hędzulek, in a similar to our age group found that the occurrence of mandibular yawning was detected in 70% of respondents. In turn, Maciejewska-Szaniec [26] is of the opinion that this symptom may concern from 8.69% to 68% of patients depending on the selection of the influenza under investigation.

However, it should be remembered that functional disorders of the masticatory system may also be accompanied by, for example, rheumatoid arthritis, limiting facial mobility / and jaw swelling, pain during abduction and palpation of the stomatognathic system muscles, Parkinson's disease, evoking mandibular shaking, difficulty in chewing, cerebral stroke causing unilateral paresis, difficulty in chewing and swallowing). [37]

However, the authors' results presented in Table 9 show the existence of a large discrepancy in the literature data on comparable age groups.

V. REFERENCES

- [1] Litko M, Kleinrok J. Dysfunkcje narządu żucia u młodocianych – przegląd piśmiennictwa. *Protet Stomatol* 2007; 2: 105-111.
- [2] Prośba-Mackiewicz M, Wytrykowska A, Mackiewicz J. Subiektywne i obiektywne objawy zaburzeń czynnościowych w układzie stomatognatycznym. *Dent Forum* 2008; 1: 17-21.
- [3] Siemińska-Piekarczyk B, Zadurska M, Biedrzycka E, Pietrzak-Bilińska B, Zwierzchowska-Walendziak H, i wsp. Etiologia i objawy kliniczne bruksizmu u dzieci i młodzieży na podstawie

- piśmiennictwa i własnych obserwacji. *Czas Stomatol* 1998;1: 47-51.
- [4] Śmierciak A. Bruksizm – definicja, diagnostyka i leczenie. *Porad Stomatol* 2007; 1: 34-41.
- [5] Panek H, Nowakowska D, Maślanka T, Bruziewicz-Miklaszewska B, Krawczykowska H, *et al.* Epidemiology of Temporomandibular Dysfunctions in young adult populations studied in Department of Prosthodontics, Silesian Piast University of Medicine in Wrocław, Poland. *Dent Med Probl* 2007; 44, 1: 55-59.
- [6] Macfarlane TV, Blinkhorn AS, Davies RM, Kincey J, Worthington HV. Oro-facial pain in the community: prevalence and associated impact. *Community Dent Oral Epidemiol* 2002; 30: 52–60.
- [7] Jancelewicz M. Dysfunkcje układu stomatognatycznego narastającym problemem współczesnej opieki zdrowotnej - przy czynny wzrostu występowania tej dysfunkcji. *Hyg*, 2010; 45 (1): 17-20.
- [8] Mankiewicz M, Panek H. Występowanie parafunkcji narządu żucia u młodocianych. *Dent Med. Probl* 2005; 42, 1: 95-101.
- [9] Prośba-Mackiewicz M, Wytrykowska A, Mackiewicz J. Subiektywne i obiektywne objawy zaburzeń czynnościowych w układzie stomatognatycznym. *Dent Forum* 2008; 1: 17-21.
- [10] Beyron H. Optimal occlusion. W: Ramfjord SP, *et al.* Symposium on Occlusion. *Dent Clin North Am* 1969; 13: 537- 554.
- [11] Rieder CE. The prevalence and magnitude of mandibular displacement in a survey population. *J Prosthet Dent* 1978; 39, 324- 329.
- [12] Reding GR, Rubright WC, Zimmerman SO. Incidence of bruxism. *J Dent Res* 1966; 45: 1198- 1204.
- [13] Solberg WK, Woo MW, Houston JG. Prevalence of mandibular dysfunction in young adults. *J Am Den. Assoc* 1979; 98: 25- 34.
- [14] Glaros A. Incidence of diurnal and nocturnal bruxism. *J Prosthet Dent* 1981; 45: 545- 549.
- [15] Gross AJ, Rivera- Morales WC, Gale EN. A prevalence study of symptoms associated with TM disorders. *J Craniomandib Disord Facial Oral Pain* 1988; 2: 191- 195.
- [16] Goulet JP, Lund JP, Lavigne G. Jaw pain: an epidemiologic survey among French Canadians in Quebec. *J Den. Res* 1992; 71: 150.
- [17] Goulet JP, *et al.* Daily clenching, nocturnal bruxism and stress and their association with TMD symptoms. *J Orofac Pain* 1993; 7: 120-124.
- [18] Shetty S, *et al.* Bruxism: a literature review. *J Indian Prosthodont Soc* 2010; 10: 141- 144.
- [19] Lobbezoo F, van der Zaag J, Naeije M. Bruxism: its multiple causes and its effects on dental implants – an updated review. *J Oral Rehab* 2006; 33: 293- 300.
- [20] Manfredini D, Lobbezoo F. Role of psychosocial factors in the etiology of bruxism. *J Orofac Pain* 2009; 23: 153- 166.
- [21] Dos Santos J. Occlusion, Principles & Treatment. London; Quintessence Pub, 2007.
- [22] Casanova-Rosado JF, Medina-Solis CE, Vallejos-Sanchez AA, Casanova-Rosado AJ, Prado BH, Ávila-Burgos L. Prevalence and associated factors for temporomandibular disorders in a group of Mexican adolescents and youth adults. *Clin Oral Invest* 2005; 11: 21-29.
- [23] Joško-Wajner N. O bruksizmie raz jeszcze. *TPS* 2009; 9: 79-81.
- [24] Śmierciak A. Bruksizm – definicja, diagnostyka i leczenie. *Porad Stomatol* 2007; 1: 34-41.
- [25] Ziółkowska-Kochan M, Kochan J, Pracka D, Drózd W, Borowska A. Bruksizm - problem interdyscyplinarny. *Czas Stomatol* 2007; 6: 391-397.
- [26] Maciejewska-Szaniec. Ocena współwystępowania zaburzeń czynnościowych układu stomatognatycznego i objawów otologicznych. Rozprawa na stopień doktora nauk medycznych. Poznań; Uniwersytet Medyczny im. K. Marcinkowskiego w Poznani, 2014.
- [27] Sójka A, Hędzalek W. Ocena występowania objawów dysfunkcji narządu żucia w grupie studentów 20-25 lat. *Protet Stomatol* 2008; LVIII, 4: 259-266.
- [28] Buergers R, Kleinjung T, Behr M, Vielsmeier V. Is there a link between tinnitus and temporomandibular disorders? *J Prosthet Dent* 2013. [online] [cited 2018 May 30] Available from: URL:<http://www.sciencedirect.com/science/article/pii/S0022391313002552>
- [29] Cooper BC, Kleinberg I: Examination of large patient population for the presence of symptoms and signs of temporomandibular disorders. *Cranio* 2007; 25: 114-119.
- [30] De Felicio CM, Faria TG, Silva MA, da Silva MA, de Aquino AM, Junqueira CA. Temporomandibular disorder: relationship between otologic and orofacial symptoms. *Rev Bras Otorinolaringol* 2004; 70 (6): 786-793.
- [31] De Felicio CM, Pimenta F CL, De Oliveira M. Otologic symptoms of temporomandibular disorder and effect of orofacial myofunctional therapy. *J Craniomandibular Pract* 2008; 4: 1-6.
- [32] Michalak M, Wysokińska-Miszczuk J, Wilczak M, Paulo M, Bozyk A, Borowicz J. Correlation between eye and ear symptoms and lack of teeth, bruxism and other parafunctions in population of 1006 patients in 2003-2008. *Arch Med Sci* 2012; 8, 1: 104-110.
- [33] Costen JB. A syndrome of ear and sinus symptoms dependent upon disturbed function of the temporomandibular joint. *Ann Otol Rhinol Laryngol* 1984; 3: 1-4.
- [34] Zabłocki S. Zespoły i objawy chorobowe w otorynolaryngologii. Gdańsk; Via Medica, 2004.
- [35] Pinto OF. A new structure related to the temporomandibular joint and middle ear. *J Pros Dent*, 1962; 12: 95-103.
- [36] Myrhaug H. The incidence of ear symptoms in cases of malocclusion and temporo-mandibular joint disturbances. *Br J Oral Surg* 1964; 2: 28-32.
- [37] Paparo F, Fatone FM, Ramieri V. Anatomic relationship between trigeminal nerve and temporomandibular joint. *Eur Rev Med Pharmacol Sci* 2008; 12: 15-18.
- [38] Nawrocka-Furmanek J, Rusiniak-Kubik K, Mierzwińska-Nastalska E, Zadurska M, Siemińska-Piekarczyk B, *i wsp.* Występowanie dysfunkcji narządu żucia w zależności od zaburzeń okluzji i wad zgryzu wśród młodych dorosłych. *Protet Stomatol* 2007; LVII, 3: 183-191.